



Key Vocabulary

ten million

millions

thousands

hundreds

tens

ones

zero

place value

greater than

less than

order

round

rounded

negative number

partition

digit

interval

sequence

linear sequence

Common Factors

Factors of 48

1 2 3 4 6 8 12 16 24 48

Factors of 30

1 2 3 5 6 10 15 30

Common factors: 1, 2, 3, 6

Primes

A prime number has only 1 and itself as factors: 2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43

A composite number has factors other than 1 and itself.

Mental Calculations and Estimation

Order of calculations:

$50 \times 34 \times 2 = 50 \times 2 \times 34 = 100 \times 34 = 3400$

Money: $\pounds 8.99 + \pounds 3.49 = \pounds 12.48$

Use $\pounds 9 + \pounds 3.50 = \pounds 12.50$ and subtract 2p

Estimate on a number line



Subdivide line to estimate: 17

Common Multiples

Multiples of 3

3 ... 18 21 24 ... 39 42

Multiples of 7

7 14 21 28 35 42

Common multiples: 21, 42...

Squares and Cubes

Square numbers result from a number being multiplied by itself (e.g. $5 \times 5 = 25$):

1, 4, 9, 16, 25, 36, 49, 64, 81, 100

Cube numbers result from a number being multiplied by itself twice ($2 \times 2 \times 2 = 8$):

1, 8, 27, 64, 125

Reason from Known Facts

$90 \div 10 = 9$ so $90 \div 20 = 4.5$ and $90 \div 5 = 18$

$16 \times 9 = 144$ so $1.6 \times 9 = 14.4$

$4352 \div 17 = 256$

so $256 \times 18 = 4352 + 256 = 4608$

$3786 + 2850 = 6636$

so $4786 + 2850 = 7636$

and $2786 + 3850 = 6636$

and $8636 - 3786 = 4850$

Long Division

		1	2	0	r	3
14	1	6	8	3		
	1	4	0	0		
		2	8	3		
		2	8	0		
				3		

Adding and Subtracting Mixed Numbers

Add or subtract the whole numbers and fractions separately.

$$2\frac{2}{5} + 1\frac{3}{10}$$

$$2+1=3$$

$$\frac{2}{5} + \frac{3}{10} = \frac{4}{10} + \frac{3}{10} = \frac{7}{10}$$

$$3 + \frac{7}{10} = 3\frac{7}{10}$$

$$2\frac{1}{2} - 1\frac{1}{4}$$

$$2-1=1$$

$$\frac{1}{2} - \frac{1}{4} = \frac{2}{4} - \frac{1}{4} = \frac{1}{4}$$

$$1 + \frac{1}{4} = 1\frac{1}{4}$$

Convert the mixed numbers to improper fractions.

$$2\frac{2}{5} = \frac{12}{5}$$

$$1\frac{3}{10} = \frac{13}{10}$$

$$2\frac{1}{2} = \frac{5}{2}$$

$$1\frac{1}{4} = \frac{5}{4}$$

$$\frac{12}{5} + \frac{13}{10} = \frac{24}{10} + \frac{13}{10} = \frac{37}{10}$$

$$\frac{5}{2} - \frac{5}{4} = \frac{10}{4} - \frac{5}{4} = \frac{5}{4}$$

$$\frac{37}{10} = 3\frac{7}{10}$$

$$\frac{5}{4} = 1\frac{1}{4}$$

Simplify Fractions

$$\frac{9}{12}$$

Factors of 9:
1, 3, 9

Factors of 12:
1, 2, 3, 4, 6, 12

$$\frac{9}{12} = \frac{3}{4}$$



Numbers to Ten Million

3 926 471

Millions	Hundred Thousands	Ten Thousands	Thousands	Hundreds	Tens	Ones
3	9	2	6	4	7	1

three million, nine hundred and twenty-six thousand, four hundred and seventy-one




3 926 471
3 926 000 471



Metric to Imperial Conversions

metric (new)	imperial (old)
2.5 centimetres	1 inch
1 kilogram	2.2 pounds
4.5 litres	1 gallon



Key Vocabulary

term to term rule

variable

unknown

expression

equation

formula

one-step equation

two-step equation

substitution





pairs of unknowns

enumerate

Linear Number Sequences

A linear number sequence is a sequence where each value increases or decreases by the same amount each time. Each number in a linear number sequence is called a **term**. The constant change between each number is called the term to term rule. To identify the **term to term rule**, find the difference between two adjacent terms.

When you know the term to term rule, you can use it to find the next number in the sequence. It can also be used to find a missing number within a sequence.



Forming Expressions

An expression is a group of numbers, letters and operation symbols.

Add 14 to a

$$a + 14$$

Subtract 20 from b

$$b - 20$$

Multiply c by 4

$$4c$$

12 more than d

$$d + 12$$

Multiply e by 3 and subtract 5

$$3e - 5$$

Add 12 to f and then multiply by 2

$$2(f + 12)$$

Forming Equations

$$a + 14 = 20$$

$$b - 20 = 15$$

$$4c = 28$$

$$d + 12 = 30$$

$$3e - 5 = 10$$

$$2(f + 12) = 44$$

An equation is a number statement with an equal sign (=). Expressions on either side of the equal sign are of equal value.

Formulas / Formulae

(The word formula has two possible plural forms, formulae and formulas.)

A formula is a special type of equation that shows the relationship between different substituted variables. Formulas are often used in geometry to find area and volume.

Area of rectangle =
length $\times$ width

Area of triangle =
(base $\times$ height) $\div$ 2

(12.5 $\times$ hours worked)
+ 25 = cost of job

Equations with Pairs of Unknowns

In an equation with two unknown numbers, there may be **several** possible values for the unknowns that will balance the equation.

$ab = 18$	
a	b
1	18
2	9
3	6
6	3
9	2
18	1


$2a + b = 10$	
a	b
2	6
3	4
4	2
5	0

Enumerating Possibilities

Enumerating means making a complete list of answers to a problem.

- Use a system for finding the possibilities.
- Organise your findings in an ordered list or table.
- Have a way of deciding when all possibilities have been found.

There are four ice cream flavours.




Two scoops of two different flavours give six possible combinations.

- chocolate and strawberry
- chocolate and vanilla
- chocolate and mint
- strawberry and vanilla
- strawberry and mint
- vanilla and mint

Solving One-Step and Two-Step Equations


In algebra, missing numbers in equations are represented by letters. Any letter can be used but often the letter x is used. An algebraic x is written to look different to a normal letter ' x ' to avoid confusion.

$$3x = 15$$



The multiplication sign is not used in algebra to avoid confusing it with the algebraic x used to show a missing number. Inverse operations are used to isolate the letter on one side of the equation.

$$2x + 4 = 10$$



Converting Units

Knowledge Organiser

Key Vocabulary

mass
gram
kilogram
capacity
volume
millilitre
litre
millimetre
centimetre
kilometre
foot
inch
ounce
pound
stone
pint
gallon

Converting Mass

$$1 \text{ tonne} = 1000\text{kg}$$


$$1000\text{g} = 1\text{kg}$$

$$\frac{1}{10} \text{ kg} = 0.1\text{kg} = 100\text{g}$$

$$\frac{1}{4} \text{ kg} = 0.25\text{kg} = 250\text{g}$$

$$\frac{1}{2} \text{ kg} = 0.5\text{kg} = 500\text{g}$$

$$\frac{3}{4} \text{ kg} = 0.75 = 750\text{g}$$



Converting Capacity

$$1000\text{ml} = 1\text{l}$$


$$\frac{1}{10} \text{ l} = 0.1\text{l} = 100\text{ml}$$

$$\frac{1}{4} \text{ l} = 0.25\text{l} = 250\text{ml}$$

$$\frac{1}{2} \text{ l} = 0.5\text{l} = 500\text{ml}$$

$$\frac{3}{4} \text{ l} = 0.75\text{l} = 750\text{ml}$$

$$\frac{1}{100} \text{ l} = 0.01\text{l} = 10\text{ml}$$



Converting Length

$$1000\text{m} = 1\text{km}$$

$$100\text{cm} = 1\text{m}$$



$$10\text{mm} = 1\text{cm}$$

$$\frac{1}{2} \text{ m} = 0.5\text{m} = 50\text{cm}$$

$$\frac{1}{4} \text{ m} = 0.25\text{m} = 25\text{cm}$$


$$\frac{3}{4} \text{ m} = 0.75\text{m} = 75\text{cm}$$

$$\frac{1}{10} \text{ m} = 0.01\text{m} = 10\text{cm}$$



Miles to Kilometres

You might measure the length of a road or the distance between two cities in miles or kilometres.



Time


Minute 1 minute = 60 seconds

Hour 1 hour = 60 minutes

Day 1 day = 24 hours

Week 1 week = 7 days

Year 1 year = 12 months = 52 weeks = 365 days



Imperial Measures


Things that could be measured using imperial units:

- Someone's height in feet and inches
- The mass of a bag of sugar in ounces
- The mass of a sack of potatoes in pounds
- A person's mass in stones
- A carton of milk in pints
- The amount of water in a bath in gallons

1 foot = 12 inches
1 pound = 16 ounces
1 stone = 14 pounds
1 gallon = 8 pints

Metric to Imperial Conversions

metric (new)	imperial (old)
2.5 centimetres	1 inch
1 kilogram	2.2 pounds
4.5 litres	1 gallon



Decimals

Knowledge Organiser

Key Vocabulary

decimal place

decimal fraction

recurring decimal

equivalent fraction

tenth

sharing

partitioning

exchanging

rounding to 3d.p.

hundredth

thousandth

equal to

remainder

grouping

Place Value

Tens	Ones	tenths	hundredths	thousandths
	1 1 1	0.1 0.1 0.1 0.1	0.01 0.01	0.001 0.001 0.001 0.001

$$3 + \frac{4}{10} + \frac{2}{100} + \frac{6}{1000}$$

3.426

$$3 + 0.4 + 0.02 + 0.006$$

1	2	3	4	5	6	7	8	9
0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.001	0.002	0.003	0.004	0.005	0.006	0.007	0.008	0.009

Fractions to Decimals

$$\frac{7}{20} = \frac{35}{100} \text{ or } 0.35$$

$$\frac{7}{25} = \frac{28}{100} \text{ or } 0.28$$

$$\frac{7}{50} = \frac{14}{100} \text{ or } 0.14$$

$$\frac{8}{200} = \frac{4}{100} \text{ or } 0.04$$

When the denominator is not a factor or multiple of 100

$$\frac{7}{8} = 7 \div 8$$

	0	8	7	5
8	7	0	0	0

Dividing Decimals by Integers

$$8.12 \div 4$$

	2	0	3
4	8	1	2

$$6.93 \div 3 = 2.31$$

Ones	tenths	hundredths
1 1	0.1 0.1 0.1	0.01
1 1	0.1 0.1 0.1	0.01
1 1	0.1 0.1 0.1	0.01

Decimals

Knowledge Organiser

Multiplying and Dividing by 10, 100 and 1000

Thousands	Hundreds	Tens	Ones	tenths	hundredths	thousandths
			2	0	8	
		$\times 10$	0	8		
		2	0	8		
			2	0	8	

Thousands	Hundreds	Tens	Ones	tenths	hundredths	thousandths
		4	3	5		
$\times 100$		4	3	5		
4	3	5	0			
		4	3	5		

Thousands	Hundreds	Tens	Ones	tenths	hundredths	thousandths
			1	3	5	1
$\times 1000$			1	3	5	1
1	3	5	1			
			1	3	5	1



Multiplying Decimals by Integers

	3	4	5
$\times$			3
1	0	3	5
	1	1	


$$3.21 \times 3 = 9.63$$

Ones	tenths	hundredths
1 1 1	0.1 0.1	0.01
1 1 1	0.1 0.1	0.01
1 1 1	0.1 0.1	0.01


Decimal Numbers as Fractions




$$\frac{1}{100} = 0.01$$




$$\frac{50}{100} = \frac{1}{2} = 0.5$$



$$\frac{25}{100} = \frac{1}{4} = 0.25$$



$$\frac{75}{100} = \frac{3}{4} = 0.75$$



$$\frac{20}{100} = \frac{1}{5} = 0.2$$

$$\frac{1}{3} = 0.33$$

$$\frac{1}{8} = 0.125$$

$$\frac{1}{1000} = 0.001$$

Four Operations

Knowledge Organiser

Key Vocabulary

Add

Total

Make

Plus

Sum

More

Altogether

Difference

Leave

Subtract

Difference between

Less

Minus

Take away

Mentally, Orally

Column Addition

Column Subtraction

Estimate

Inverse operation

Solve problems

Number facts

Place Value

Complex

Add and Subtract Whole Numbers

Column Method

	4	5	8	6	4
+	2	3	4	9	7
	6	9	3	6	1
		1	1	1	

Starting with the ones, add each column in turn.
Regroup tens, hundreds, thousands, ten thousands as required.

	3	5	6	13	1
-		3	4	7	6
	3	2	2	6	6

Starting with the ones, subtract each column in turn.
Exchange tens, hundreds, thousands and/or ten thousands as required.

Multiply up to 4-digit by 2-digit

1	3	2	
	1	5	4
×		2	6
	9	2	4
3	0	8	0
4	0	0	4
1	1		

Start with the ones.

$$154 \times 6 = 924$$

$$154 \times 20 = 3080$$

$$3080 + 924 = 4004$$

Order of Operations

B	Brackets	$10 \times (4 + 2) = 10 \times 6 = 60$
O	Order	$5 + 2^2 = 5 + 4 = 9$
D	Division	$10 + 6 \div 2 = 10 + 3 = 13$
M	Multiplication	$10 - 4 \times 2 = 10 - 8 = 2$
A	Addition	$10 \times 4 + 7 = 40 + 7 = 47$
S	Subtraction	$10 \div 2 - 3 = 5 - 3 = 2$

Four Operations

Short Division

Start from the left.

		4	4	0	• 5	
12	5	⁵ 2	⁴ 8	6	⁶ 0	

$5 \div 12 = 0 \text{ r}5$
 $52 \div 12 = 4 \text{ r}4$
 $48 \div 12 = 4$
 $6 \div 12 = 0 \text{ r}6$

Long Division

		1	2	0	r	3
14	1	6	8	3		
	1	4	0	0		
		2	8	3		
		2	8	0		
				3		

Common Factors

Factors of 48

1	2	3	4	6	8	12	16	24	48
---	---	---	---	---	---	----	----	----	----

Factors of 30

1	2	3	5	6	10	15	30
---	---	---	---	---	----	----	----

Common factors: 1, 2, 3, 6

Primes

A prime number has only 1 and itself as factors: 2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43

A composite number has factors other than 1 and itself.

Mental Calculations and Estimation

Order of calculations:

$$50 \times 34 \times 2 = 50 \times 2 \times 34 = 100 \times 34 = 3400$$

Money: $\pounds 8.99 + \pounds 3.49 = \pounds 12.48$

Use $\pounds 9 + \pounds 3.50 = \pounds 12.50$ and subtract 2p

Estimate on a number line



Subdivide line to estimate: 17

Knowledge Organiser

Common Multiples

Multiples of 3

3	...	18	21	24	...	39	42
---	-----	----	----	----	-----	----	----

Multiples of 7

7	14	21	28	35	42
---	----	----	----	----	----

Common multiples: 21, 42...

Squares and Cubes

Square numbers result from a number being multiplied by itself (e.g. $5 \times 5 = 25$):

1, 4, 9, 16, 25, 36, 49, 64, 81, 100

Cube numbers result from a number being multiplied by itself twice ($2 \times 2 \times 2 = 8$):

1, 8, 27, 64, 125

Reason from Known Facts

$90 \div 10 = 9$ so $90 \div 20 = 4.5$ and $90 \div 5 = 18$

$16 \times 9 = 144$ so $1.6 \times 9 = 14.4$

$4352 \div 17 = 256$

so $256 \times 18 = 4352 + 256 = 4608$

$3786 + 2850 = 6636$

so $4786 + 2850 = 7636$

and $2786 + 3850 = 6636$

and $8636 - 3786 = 4850$

Fractions

Knowledge Organiser

Key Vocabulary

numerator

denominator

proper fraction

improper fraction

factor

highest common multiple

lowest common multiple

equivalents

common numerator

common denominator

decimal equivalent

simplify

simplest form


mixed number

whole number

mixed number



Simplify Fractions




Factors of 9:

1, 3, 9

Factors of 12:

1, 2, 3, 4, 6, 12




Compare and Order Fractions

Use the Common Denominator





Multiples of 5:

5, 10, **15**




Multiples of 3:

3, 6, 9, 12, **15**




Use the Common Numerator







Multiples of 5:

5, **10**, 15



Multiples of 10:

10, 20



Fractions

Knowledge Organiser

Adding and Subtracting Proper Fractions

Same Denominators



$$\frac{4}{7} + \frac{2}{7} = \frac{6}{7}$$



$$\frac{8}{11} - \frac{3}{11} = \frac{5}{11}$$

Different Denominators

$$\frac{2}{7} + \frac{3}{5}$$

Multiples of 7: 7, 14, 21, 28, **35**

Multiples of 5: 5, 10, 15, 20, 25, 30, **35**

$$\frac{2}{7} = \frac{10}{35}, \frac{3}{5} = \frac{21}{35}$$

$$\frac{10}{35} + \frac{21}{35} = \frac{31}{35}$$

$$\frac{9}{10} - \frac{1}{4}$$

Multiples of 10: 10, **20**

Multiples of 4: 4, 8, 12, 16, **20**

$$\frac{9}{10} = \frac{18}{20}, \frac{1}{4} = \frac{5}{20}$$

$$\frac{18}{20} - \frac{5}{20} = \frac{13}{20}$$

Multiplying Proper Fractions

Multiplying Fractions by Fractions

$$\frac{1}{2} \times \frac{1}{3} = \frac{1}{2} \times \frac{1}{3} = \frac{1}{6}$$

Multiplying Fractions by Whole Numbers



$$\frac{2}{5} \times 3 \rightarrow$$


$$3 = \frac{3}{1}$$

$$\frac{2}{5} \times \frac{3}{1} = \frac{6}{5} = 1 \frac{1}{5}$$

Adding and Subtracting Mixed Numbers

Add or subtract the whole numbers and fractions separately.

$$2 \frac{2}{5} + 1 \frac{3}{10}$$

$$2 + 1 = 3$$

$$\frac{2}{5} + \frac{3}{10} = \frac{4}{10} + \frac{3}{10} = \frac{7}{10}$$

$$3 + \frac{7}{10} = 3 \frac{7}{10}$$

$$2 \frac{1}{2} - 1 \frac{1}{4}$$

$$2 - 1 = 1$$

$$\frac{1}{2} - \frac{1}{4} = \frac{2}{4} - \frac{1}{4} = \frac{1}{4}$$

$$1 + \frac{1}{4} = 1 \frac{1}{4}$$

Convert the mixed numbers to improper fractions.

$$2 \frac{2}{5} + 1 \frac{3}{10}$$

$$2 \frac{1}{2} - 1 \frac{1}{4}$$

$$2 \frac{2}{5} = \frac{12}{5}$$

$$1 \frac{3}{10} = \frac{13}{10}$$

$$2 \frac{1}{2} = \frac{5}{2}$$

$$1 \frac{1}{4} = \frac{5}{4}$$

$$\frac{12}{5} + \frac{13}{10} = \frac{24}{10} + \frac{13}{10} = \frac{37}{10}$$

$$\frac{37}{10} = 3 \frac{7}{10}$$

$$\frac{5}{2} - \frac{5}{4} = \frac{10}{4} - \frac{5}{4} = \frac{5}{4}$$

$$\frac{5}{4} = 1 \frac{1}{4}$$

Dividing Fractions by Whole Numbers

$$\frac{2}{5} \div 2 = \frac{1}{5}$$

Multiplication and division are the inverse of one another so:

$\div 2$ is the same as $\times \frac{1}{2}$

$$\frac{2}{5} \times \frac{1}{2} = \frac{2}{10}$$

Number and Place Value

Knowledge Organiser

Key Vocabulary

Compare and Order

ten million

millions

thousands

hundreds

tens

ones

zero

place value

greater than

less than

order

round

rounded

negative number

partition

digit

interval

sequence

linear sequence

equals

$$26 + 38 = 8 \times 8$$

Both calculations have the value 64.

greater than

$$223\,873 > 98\,256$$

The number on the left has 2 hundred thousands and the number on the right has 0 hundred thousands.

less than

$$901\,198 < 1\,091\,098$$

The number on the right has 1 million and the number on the left has 0 millions.

smallest

81 782

127 352

127 835

137 019



200 002

greatest

Negative Numbers

$$3 - 8 = -5$$

$$-6 + 11 = 5$$




Numbers to Ten Million


3 926 471

Millions	Hundred Thousands	Ten Thousands	Thousands	Hundreds	Tens	Ones
3	9	2	6	4	7	1

three million, nine hundred and twenty-six thousand, four hundred and seventy-one




3 926 471
3 926 000 471




Round Any Number

Rounding to the nearest 1000




Rounding to the nearest 10 000




visit [twinkl.com](https://www.twinkl.com)

Rounding to the nearest 100 000



Rounding to the nearest 1 000 000



Percentages

Knowledge Organiser

Key Vocabulary

per cent (%) =
'out of 100'

percentage

discount

equivalent fraction

equivalent decimal

convert


compare

order


the whole




Equivalent Fractions, Decimals and Percentages




$$\frac{50}{100} = \frac{1}{2} = 0.5 = 50\%$$




$$\frac{25}{100} = \frac{1}{4} = 0.25 = 25\%$$




$$\frac{10}{100} = \frac{1}{10} = 0.1 = 10\%$$



$$\frac{75}{100} = \frac{3}{4} = 0.75 = 75\%$$



$$\frac{1}{100} = 0.01 = 1\%$$



$$\frac{20}{100} = \frac{2}{10} = 0.2 = 20\%$$


Fractions to Percentages

$$\begin{array}{c} \times 2 \\ \swarrow \quad \searrow \\ \frac{15}{50} = \frac{30}{100} = 0.3 = 30\% \\ \nearrow \quad \nwarrow \\ \times 2 \end{array}$$


$$\begin{array}{c} \div 2 \\ \swarrow \quad \searrow \\ \frac{60}{200} = \frac{30}{100} = 0.3 = 30\% \\ \nearrow \quad \nwarrow \\ \div 2 \end{array}$$

Order Fractions, Decimals and Percentages


$$\frac{3}{10} > 25\% > 0.2$$



$$\frac{30}{100} = 30\%$$




$$\frac{25}{100} = 25\%$$




$$\frac{20}{100} = 20\%$$


$$80\% = 0.8 = \frac{4}{5}$$



$$\frac{80}{100} = 80\%$$



$$\frac{80}{100} = 80\%$$



$$\frac{80}{100} = 80\%$$





Finding a Percentage of an Amount

$$50\% = \frac{1}{2} \text{ so we can divide by 2}$$

$$10\% = \frac{1}{10} \text{ so we can divide by 10}$$


$$25\% = \frac{1}{4} \text{ so we can divide by 4}$$

$$1\% = \frac{1}{100} \text{ so we can divide by 100}$$



Percentages – Missing Values

Whole value (100%) of bar model = ?



We know 10% = 15 $10\% \times 10 = 100\%$ (the whole) so $15 \times 10 = 150$

Key Vocabulary

perimeter

area

volume

cubic units (e.g. cm^3)

cuboid

width

length

rectangle


rectilinear

parallelogram

perpendicular height

Area of Rectangles

length $\times$ width = area of a rectangle




Counting squares:

area = 18cm^2

Use formula:

$6\text{cm} \times 3\text{cm}$

area = 18cm^2




$8\text{cm} \times 4\text{cm}$ area = 32cm^2

Perimeter of Rectangles

perimeter = length + width + length + width or (length + width) $\times$ 2




$5\text{cm} + 4\text{cm} + 5\text{cm} + 4\text{cm}$
area = 18cm^2



$(6 + 2) \times 2$
area = 16cm^2

Area of Triangles



base $\times$ perpendicular height $\div$ 2 = area of a triangle



$8\text{cm} \times 3\text{cm} \div 2$
area = 12cm^2

perpendicular height = 5cm

$6\text{cm} \times 5\text{cm} \div 2$
area = 15cm^2



Counting squares:

6 whole squares = 6cm^2

6 half squares = 3cm^2

$6\text{cm}^2 + 3\text{cm}^2 = 9\text{cm}^2$

area = 9cm^2


Using formula:

$6\text{cm} \times 3\text{cm}$


$\div 2 = 9\text{cm}^2$

Perimeter and Area

Shapes with the same area can have different perimeters.




area = 8cm^2 perimeter = 12cm




area = 8cm^2 perimeter = 18cm

Shapes with the same perimeter can have different areas.



area = 8cm^2 perimeter = 12cm




area = 5cm^2 perimeter = 12cm


Volume - Counting Cubes



= 1cm^3



11cm^3




27cm^3

Area of Parallelograms


base $\times$ perpendicular height = area of a parallelogram

A parallelogram can be transformed into a rectangle.



Volume of Cuboids

length $\times$ width $\times$ height = volume of a cuboid



Multiply dimensions in **any** order:

$3\text{cm} \times 6\text{cm} \times 4\text{cm}$

volume = 72cm^3

Key Vocabulary

translate

translation

reflect

reflection

up

down

right

left

coordinates

quadrant

x-axis


y-axis

horizontal

vertical

Four Quadrants

Coordinates can use positive and negative numbers. Whether positive or negative, the x-axis coordinate is written first, followed by the y-axis coordinate.




Look at the circle. It is 3 units along the x-axis and 4 down the y-axis. Its coordinates are (3,-4).

Completing Shapes

Using the properties of a shape, a polygon can be completed on a grid.


To make a square, think of the square's properties.



All of a square's sides are the same length. If the completed sides are 2 units in length, the missing point must complete two more sides of 2 units.

To make a right-angled triangle, think of the triangle's properties.

A right-angled triangle should have three sides with one 90° angle.




Translation

A shape is translated when it is moved without being rotated or resized. Every point of the shape moves the same distance and in the same direction.




Shape 1 has been translated 4 units left and 3 units down.




Reflections

A shape is reflected when it is flipped over a line which acts as a mirror. Every point on the original shape is the same distance from the mirror line as the same point on the reflected shape. The original triangle has been reflected in the x-axis and in the y-axis.



Missing Coordinates

Shapes can be shown on unmarked grids.



Point a is in the same position along the x-axis as (5,2) and in the same position on the y-axis as (7,9).

Point a (5,9)

Point b is in the same position on the y-axis as (10,4). Both triangles will have the same width. The width of the right-hand triangle is 3. This means that the width of the left-hand triangle is also 3.

Point b (2,4)


Properties of Shapes

Knowledge Organiser

Key Vocabulary

angle
 right angle
 acute
 obtuse
 reflex
 protractor
 horizontal
 vertical
 parallel
 perpendicular
 polygon
 regular
 irregular
 two-dimensional
 three-dimensional
 flat face
 curved surface
 edge
 curved edge
 vertex
 vertices
 apex
 radius
 diameter
 circumference

Angle Types



Acute Angles

Any angle that measures less than 90° is called an **acute** angle.



Obtuse Angles

Any angle that measures greater than 90° and less than 180° is called an **obtuse** angle.




Reflex Angles

Any angle that measures greater than 180° is called a **reflex** angle.


Calculating Angles




Angles on a straight line always total 180° .



Angles around a point always total 360° .



Opposite angles that share a vertex are equal.




$\frac{1}{4}$ turn
 90°




$\frac{1}{2}$ turn
 180°



$\frac{3}{4}$ turn
 270°




1 turn
 360°




Multiples of 90° can be used as descriptions of a turn.

Angles in a Triangle



$$a + b + c = 180^\circ$$

Angles in a Quadrilateral



$$a + b + c + d = 360^\circ$$


Properties of Shapes

Using a Protractor

Place the cross or circle at the point of the angle you are measuring.

Read from the zero on the outer scale of your protractor.

Count the degree lines carefully.




Parts of Circles


A circle is a 2D shape. The perimeter of a circle is called the **circumference** (c). The distance across the circle, passing through the centre, is called the **diameter** (d).

The distance from the centre of the circle to the circumference is called the **radius** (r).

$$r \times 2 = d \quad \frac{d}{2} = r$$



Nets of 3D Shapes



A shape net shows which 2D shapes can be folded and joined to make a 3D shape. When you are drawing a net, or solving a problem involving a shape net, think carefully about where the edges of the faces meet.




Knowledge Organiser

Angles in Regular Polygons

As the number of sides of a polygon increases by one, the total of the interior angles increases by 180° . When n = number of sides, this formula can be used to find the size of each angle in a **regular polygon**:

$$\text{Sum of Interior Angles} = (n - 2) \times 180^\circ$$

$$\text{Each Angle} = \frac{(n - 2) \times 180^\circ}{n}$$



Pentagon

$$n = 5$$

$$(5 - 2) \times 180^\circ = 540^\circ$$

$$540^\circ \div 5 = 108^\circ$$



Hexagon

$$n = 6$$

$$(6 - 2) \times 180^\circ = 720^\circ$$

$$720^\circ \div 6 = 120^\circ$$

Properties of 3D Shapes

3D shapes have three dimensions – **length**, **width** and **depth**.

A **polyhedron** is a 3D shape with flat faces. Spheres, cylinders and cones are not polyhedrons as they have curved surfaces.

Cube  6 square faces 12 edges 8 vertices	Tetrahedron  4 triangular faces 6 edges 4 vertices	Sphere  1 curved surface 0 edges 0 vertices
Cuboid  6 faces 12 edges 8 vertices	Octahedron  8 faces 12 edges 6 vertices	Triangular prism  5 faces 9 edges 6 vertices
Square-based pyramid  5 faces 8 edges 5 vertices	Cone  1 circular face 1 curved surface 1 curved edge 1 apex	Cylinder  2 circular faces 1 curved surface 2 curved edges 0 vertices